

## **“Simulation of Extreme Arctic Cyclones in IPCC AR5 Experiments”**

PI: Stephen Vavrus  
Nelson Institute Center for Climatic Research  
1225 W. Dayton Street  
Madison, WI 53706  
phone: 608-265-5279, fax: 608-263-4190, email: [sjvavrus@wisc.edu](mailto:sjvavrus@wisc.edu)

Award Number: N000141110611

### **LONG-TERM GOALS**

The primary goals of this project are to assess the ability of the current generation of global climate models (GCMs) to simulate extreme Arctic cyclones and identify changes in the characteristics of these storms caused by greenhouse-forced climate change to present.

### **OBJECTIVES**

These goals are being addressed through the following questions. First, how realistically does the widely used CCSM4 GCM simulate the observed characteristics of extreme Arctic cyclones and how sensitive are these events to the horizontal resolution of the model? Second, do other GCMs generate such storms and, if so, are there any common characteristics among models that successfully do so? Third, does the preferred location of these systems and their impacts shift as the cyclogenetic baroclinic zone induced by the sea ice margin migrates poleward with time? Fourth, what do these state-of-the-art climate models suggest about changes in the frequency vs. intensity of extreme Arctic cyclones?

### **APPROACH**

I am targeting these objectives through a retrospective analysis of the transient 20th century simulations (spanning years 1850-2005) among the GCMs participating in the latest Coupled Model Intercomparison Project (CMIP5). I am including 14 GCMs with widely varying horizontal and vertical resolutions. These simulations are being compared with a new atmospheric reanalysis data set covering almost this entire period (1871-2008) from NOAA's Earth System Research Laboratory: the 20th Century Reanalysis, version 2, available at [www.esrl.noaa.gov/psd/data/gridded/data.20thC\\_ReanV2](http://www.esrl.noaa.gov/psd/data/gridded/data.20thC_ReanV2). This data set is described in detail by Compo et al. (2010) and provides various atmospheric fields, including sea level pressure (SLP), on daily and sub-daily time scales at 2° horizontal resolution. A higher-resolution and more reliable data set for the more recent 1979- period, NASA's Modern Era Retrospective-analysis for Research and Applications (MERRA), is also being used to evaluate the climate models. All of this work is being conducted by the PI.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>2012</b>		2. REPORT TYPE <b>N/A</b>		3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>Simulation of Extreme Arctic Cyclones in IPCC AR5 Experiments</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Nelson Institute Center for Climatic Research 1225 W. Dayton Street Madison, WI 53706</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>The original document contains color images.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>SAR</b>	18. NUMBER OF PAGES <b>8</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## WORK COMPLETED

I have mostly completed the 1850-2005 analysis of CCSM4, the primary GCM in this analysis, and its comparison with the 20<sup>th</sup> Century Reanalysis and MERRA. For assessing these changes during this period of modest greenhouse forcing, I have also made some comparisons with the model's response to much stronger greenhouse forcing during its 21<sup>st</sup>-century simulation. A more substantial portion of the project in the past year has been a retrospective, multi-model analysis of 14 GCMs in the CMIP5 data set over the 1850-2005 period. This work was presented at the Summer 2012 Community Earth System Modeling Workshop in Breckenridge, CO, and will be featured again at the upcoming 2012 AGU Fall Meeting ("Simulation of Extreme Arctic Cyclones in IPCC AR5 Experiments"). I am in the process of writing up the results for an article in Geophysical Research Letters. I also provided expert assessment of this summer's Arctic "megacyclone", which made international news as a possible expression of regional climate change and a contributor to the record low sea ice extent this month. <http://earthobservatory.nasa.gov/IOTD/view.php?id=78808>

## RESULTS

1. CCSM4 generates a significant increase in the strength and frequency of extreme Arctic cyclones in its 21<sup>st</sup> century simulation (2005-2100), with a maximum signal during autumn (Vavrus et al., 2011).
2. However, this same model does not show a strengthening of the most extreme Arctic cyclones during the 1850-2005 period, nor a significant change in their frequency.
3. The 20<sup>th</sup> Century Reanalysis does indicate a strengthening of the most extreme Arctic cyclones during similar time period (1871-2008), but most of this change occurs before 1920, when the data quality is questionable. The same conclusion generally holds for the reanalyzed frequencies of extreme cyclones, but this data set suggests noticeably higher occurrences in the high Arctic (poleward of 70°N) during the past 25 years.
4. CCSM4 produces more extreme polar cyclones than the Reanalysis, consistent with its bias toward low sea level pressure in the Arctic during almost every month (de Boer et al., 2011).
5. The CCSM4 simulations show an interesting shift in the location of extreme Arctic cyclones as a function of greenhouse warming (Figure 1). By the late 21<sup>st</sup> century, the model indicates a shift in the path of these storms, such that they emanate into the central Arctic Ocean from their preferred track over the North Atlantic-Barents Sea. However, this change is not effected by the modest greenhouse warming between 1850-2000, when the sea ice margin is further south and may act as a barrier for the production or migration of such extreme events.
6. On the whole, the GCMs in the CMIP5 collection do a reasonable job of reproducing the location, frequency, and intensity of the strongest cyclones (top 1%) in the Arctic during the recent past (1979-2005). The favored regions for these storms are within the climatological Icelandic and Aleutian Lows, and extreme cyclones reaching the Arctic Ocean occur almost exclusively as extensions from the Icelandic Low storm track, rather than from the Aleutian Low track. Simulated and observed intense Arctic cyclones are primarily marine phenomena, such that maximum frequencies occur over water, especially ice-free seas.
7. The models also correctly simulate that the most intense Arctic cyclones are primarily a wintertime phenomenon and almost never occur during summer, but the CMIP5 models have a bias of producing too many such storms during the late winter and early spring.

8. The CMIP5 models simulate a decrease in mean annual Arctic sea level pressure since 1850, consistent with expectations from greenhouse forcing, but the trend in extreme cyclone frequency is most pronounced in the vicinity of the Aleutian and Icelandic Lows, rather than over the Arctic Ocean.
9. Over the Arctic, I find no significant correlation between horizontal or vertical model resolution and simulated annual-mean SLP, mean minimum daily SLP, absolute minimum SLP, and extreme cyclone frequency. Apparently, sea level pressure biases in the models and the simulation of intense storms are affected more by internal physical processes, rather than resolution.

## IMPACTS


These results have implications for navigation, economic activities, military operations, and coastal erosion. Existing Naval operations in the Arctic Ocean and adjacent seas can be adversely affected by the passage of extreme cyclones, and future operations will likely become even more sensitive to these weather systems if reduced ice cover promotes more Naval activities and enhanced storminess in the region. A recent paper by Overeem et al. (2011) documents accelerating rates of coastal erosion in permafrost bluffs along the Beaufort Sea, due to storm-generated waves coinciding with expanding areas of open water. Presumably, this erosion will worsen with time as sea ice recedes and storms intensify.

## RELATED PROJECTS

This topic is directly related to follow-up research I recently proposed to NSF in collaboration with scientists at the National Center for Atmospheric Research (NCAR), University of Colorado, and University of Oklahoma to investigate the predictability of Arctic climate on seasonal to decadal timescales. [“Arctic Atmosphere and Sea Ice Predictability and the Interrelation on Decadal and Sub-decadal Timescales”, PIs: Steve Vavrus, Marika Holland, Jennifer Kay, Amy Solomon, Gijs de Boer, and Steven Caballa, submitted to NSF on May 10]


## REFERENCES

- Compo, G.P., J.S. Whitaker, P.D. Sardeshmukh, N. Matsui, R.J. Allan, X. Yin, B.E. Gleason, R.S. Vose, G. Rutledge, P. Bessemoulin, S. Brönnimann, M. Brunet, R.I. Crouthamel, A.N. Grant, P.Y. Groisman, P.D. Jones, M. Kruk, A.C. Kruger, G.J. Marshall, M. Maugeri, H.Y. Mok, Ø. Nordli, T.F. Ross, R.M. Trigo, X.L. Wang, S.D. Woodruff, and S.J. Worley, 2011: [The Twentieth Century Reanalysis Project](#). Quarterly J. Roy. Meteorol. Soc., 137, 1-28. DOI: 10.1002/qj.776.
- de Boer, G., W. Chapman, J. Kay, B. Medeiros, M. D. Shupe, S. Vavrus, and J. Walsh, 2011: A characterization of the present-day Arctic atmosphere in CCSM4. *J. Climate*, 2676-2695.
- Overeem, I., R. S. Anderson, C. W. Wobus, G. D. Clow, F. E. Urban, and N. Matell, 2011: Sea ice loss enhances wave action at the Arctic coast. *Geophys. Res. Lett.*, 38, L17503, doi:10.1029/2011GL048681.
- Vavrus, S., M. M. Holland, A. Jahn, D. A. Bailey, and B. A. Blazey, 2011: 21<sup>st</sup>-century Arctic climate change in CCSM4. *J. Climate*, 25, 2696-2710.




***Figure 1. Changes in the simulated frequency (events per 20-year interval) of extreme Arctic cyclones (< 950 hPa) in the CCSM4 climate model during the middle 19<sup>th</sup> century, late 20<sup>th</sup> century, early 21<sup>st</sup> century, and late 21<sup>st</sup> century.***

## Intensity of Strongest 1% of Arctic Cyclones (1850-2005)




*Figure 2. Simulated intensity (hPa) of the top percentile of Arctic cyclones in the CMIP5 models (1850-2005) compared with MERRA reanalysis (1979-2005).*

Model Average Extreme Cyclone Frequency  
Annual Means 1850–2005





***Figure 3. CMIP5-average annual frequency of extreme Arctic cyclones (deepest 1% of daily minimum SLP poleward of 50°N).***



*Figure 4. Annual cycle of extreme Arctic cyclones, as simulated by the CMIP5 models (black) and MERRA reanalysis (red).*



Model Average Trend in Extreme Cyclone Frequency  
( $\times 10^4$ ) 1850–2005



***Figure 5. Multi-model trend ( $\times 10^4$ ) among CMIP5 GCMs in extreme Arctic cyclone frequency between 1850-2005.***